

Freshwater ecosystem services resilience in a changing world

Ana Paula Portela 🕩

1. CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

2. Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, Rua do Campo Alegre s/n, 4169–007 Porto, Portugal

3. BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661 Vairão, Portugal

4. CE3C - Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

* Corresponding author: apportela@fc.ul.pt

Received: 31/01/25

Accepted: 08/05/25

Available online: 12/06/25

ABSTRACT

Freshwater ecosystem services resilience in a changing world

Healthy freshwater ecosystems provide essential ecosystem services to society such as clean water. However, freshwater ecosystems are degraded, and freshwater biodiversity is severely threatened due to anthropogenic impacts and stressors. Climate change interacts with existing stressors and may compromise the resilience of freshwater ecosystems and their services in the future. Here the aim is to review advances in assessing freshwater ecosystem services and their resilience to environmental change. This work reviews the ecosystem services provided by freshwaters, the conceptual background on ecological resilience, and examples on the resilience of freshwater ecosystems and their services. Examples from African lakes, the Pantanal wetland in Brazil and the Murray-Darling Basin riparian forests in Australia are used to understand the resilience of freshwater ecosystems and their services. This work illustrates the diverse responses of freshwater socio-ecological systems to environmental change and highlights examples of declining resilience of freshwater ecosystems and their services. However, a high degree of uncertainty still surrounds the identification of regime shifts and future ecosystem trajectories. Research is needed to understand the dynamics of freshwater socio-ecological systems and ensure resilient ecosystems and societies.

KEY WORDS: biodiversity, nature's contributions to people, rivers, lakes, and riparian ecosystems

RESUMO

Resiliência dos serviços de ecossistema aquáticos num mundo em alteração.

Os ecossistemas de águas interiores saudáveis prestam serviços de ecossistema essenciais à sociedade, como a água potável. No entanto, estes ecossistemas encontram-se degradados e a biodiversidade dos mesmos ameaçada devido a impactos antropogénicos. As alterações climáticas interagem com as pressões existentes e podem comprometer a resiliência dos ecossistemas de águas interiores e dos seus serviços no futuro. O objetivo é analisar os avanços na avaliação dos serviços de ecossistema de água doce e a sua resiliência às

alterações ambientais. Neste trabalho apresenta-se uma revisão dos serviços de ecossistema prestados pelas águas interiores, o enquadramento conceptual da resiliência ecológica e exemplos sobre a resiliência dos ecossistemas de água doce e dos seus serviços. Os exemplos de lagos africanos, da zona húmida do Pantanal no Brasil e das florestas ripícolas da bacia do rio Murray-Darling na Austrália são utilizados para compreender a resiliência dos serviços dos ecossistemas de água doce às alterações e perturbações climáticas recentes e em curso. Este trabalho ilustra a diversidade de respostas dos sistemas socio-ecológicos de água doce às alterações ambientais, realçando exemplos preocupantes de declínio da resiliência dos ecossistemas de água doce devido às alterações climáticas e por eventos extremos. No entanto, subsiste um elevado grau de incerteza quanto às mudanças de estado destes ecossistemas e às suas trajetórias futuras. A investigação da resiliência dos sistemas socio-ecológicos associados às águas interiores é necessária para compreender as suas dinâmicas e garantir ecossistemas e sociedades resilientes no futuro.

PALAVRAS CHAVE: biodiversidade, contribuição da natureza para as pessoas, rios, lagos, ecossistemas ripícolas.

This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0) License.

INTRODUCTION

Healthy freshwater ecosystems provide essential ecosystem services to society, including vital resources such as clean water. However, freshwater ecosystems are degraded, freshwater biodiversity is threatened, and an estimated 4.4 billion people globally lack access to safe drinking water (Greenwood et al., 2024). Rapid global changes including a growing human population, increasing demand for natural resources, urbanization, and intensification of agriculture have led to a focus on the exploitation of provisioning services, often at the expense of other essential services, particularly regulating services (Brauman et al., 2020). Climate change is already interacting with anthropogenic pressures and stressors, increasing the likelihood of large, potentially irreversible ecosystem changes that can have significant impacts on ecosystem services and human well-being (Forzieri et al., 2022). Thus, one of the major challenges of sustainability in the 21st century is to protect and restore ecosystems to ensure the flow of ecosystem services to society, both today and in the future (United Nations, 2022). This challenge is perhaps nowhere more pressing than in freshwater ecosystems. The strong interdependence and feedback between natural and human systems within freshwaters make them crucial social-ecological systems (Dunham et al., 2018). The ecosystem services linked to freshwater ecosystems include basic human needs and underpin social and economic well-being (Falkenmark & Wang-Erlandsson, 2021). Therefore, ensuring the resilience of freshwater ecosystem services is of significant societal and policy interest.

This work aims to provide an overview of the ecosystem services supplied by freshwaters, their resilience under global change and identify implications for management. First, I focus on the definition of ecosystem services, provide an overview of freshwater ecosystem services and discuss current limitations in their assessment. Second, I discuss the definition of resilience, the mechanisms that underpin it, and evaluate the resilience of freshwater ecosystem services to environmental changes using examples from African lakes, the Pantanal wetland in Brazil and the Murray-Darling Basin riparian forests in Australia. Third, I discuss implications for the resilience of freshwater socio-ecological systems under global change.

I. FRESHWATER ECOSYSTEM SERVICES

Definition of ecosystem services

The ecosystem services concept has been crucial in mainstreaming the link between ecosystems and human societies in research and policy. The concept and its associated theoretical framework have evolved greatly since it was first coined (Ehrlich & Ehrlich, 1981), particularly in two critical moments associated with the development of international science-policy interfaces. The first was the Millennium Ecosystem Assessment, supported by the United Nations and the Convention on Biological Diversity which mainstreamed the concept of ecosystem services leading to an exponential growth of the literature (Braat & de Groot, 2012; Gómez-Baggethun et al., 2010). The second was the establishment of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) in 2012 to create a science-policy platform for biodiversity comparable to the Intergovernmental Panel on Climate Change (Díaz et al., 2015). The Millennium Ecosystem Assessment defined ecosystem services simply as the benefits people obtain from ecosystems (Millennium Ecosystem Assessment, 2005). IPBES expanded this concept into Nature's contributions to people defined as all the contributions, positive and negative, of living nature (diversity of organisms, ecosystems, and their ecological and evolutionary processes) to people's quality of life (Díaz et al., 2018). The change aimed at addressing criticisms faced by the ecosystem services concept related to the lack of recognition of diverse values of nature, the role of culture in shaping relationships between people and nature and concerns about nature's commodification (Díaz et al., 2018). The introduction of nature's contributions to people initiated an intense debate about its novelty and value within the ecosystem services research and practice community (Kadykalo et al., 2019). It is beyond the scope of this work to address this debate. Here, this is considered a natural evolution of a field aiming to understand relationships between nature and people. The term ecosystem service is used for consistency with a large part of the referenced literature.

Freshwater ecosystem services

Freshwater ecosystem services are all those services and benefits generated by freshwater ecosystems and the interactions of land and water in ecosystems such as forests, agricultural lands, riparian areas, wetlands and water bodies (Grizzetti et al., 2016). The definition is much broader than that of hydrologic services, which focuses on the way terrestrial ecosystems affect freshwater resources through ecohydrological processes in landscapes (Brauman, 2015). By recognizing the interactions between land and water and the processes within freshwater ecosystems, the definition of freshwater ecosystem services can better reflect the ecosystem functions and services that are relevant to freshwater management.

The ecosystem services and benefits provid-

ed by freshwater ecosystems span all categories including provisioning or material contributions, regulating, and cultural or non-material contributions (Table 1; Aylward et al., 2005; Díaz et al., 2018). The initial MEA and the national assessments focused solely on hydrological services or were based on expert knowledge. Recently, there has been an increasing interest in the freshwater scientific community to contribute to the area of ecosystem services mainly by reviewing and updating the knowledge for some freshwater organisms including riparian vegetation (Riis et al., 2020), macrophytes (Thomaz, 2023), freshwater bivalves (Zieritz et al., 2022), aquatic fungi (Seena et al., 2022), freshwater fish (Holmlund & Hammer, 1999) and freshwater ecosystems namely lakes (Heino et al., 2021; Sterner et al., 2020), intermittent rivers (Datry et al., 2018; Pastor et al., 2022) and dry rivers (Nicolás Ruiz et al., 2021), small streams (Ferreira et al., 2022), wetlands (Xu et al., 2020).

Difficulties in assessment

The concept of ecosystem services has gained traction in freshwater science as evidenced by the reviews and studies published recently. However, the actual assessment of the supply and demand of freshwater ecosystem services provided by these ecosystems remains challenging. In previous studies on ecosystem services, less than 5% of indicators on supply and less than 1% of monetary valuation estimates were related to freshwater ecosystems (Egoh et al., 2012; Maes et al., 2012; Van der Ploeg et al., 2010). In studies specifically quantifying river ecosystem services, three or fewer ecosystem services were evaluated on average (Hanna et al., 2018). The most frequently assessed ecosystem services are recreation and tourism, water supply, water quality, habitat provision and erosion prevention (Hanna et al., 2018), however overall, the regulation and provision services were assessed more frequently. These trends reflect the lack of development of ecosystem services methods specifically targeting freshwater ecosystems (Riis et al., 2020). A review on riparian vegetation notes that the assessment of ecosystem services provided by different vegetation types remains based on expert

for nutritional purposes es and other materials from wild es for direct use or processing luding genetic materials) I plants (terrestrial and aquatic, eding fungi, algae) used as a source pregy plants (terrestrial and aquatic, ding fungi, algae) used for nutrition rer and lower plants (whole or- sms) used to breed new strains or etics hwater surface water used as an Hyc
up and second as a source regy to the second and a guartic, Edit the lants (terrestrial and aquatic, Edit ading fungi, algae) used for nutrition the rand lower plants (whole orsms) used to breed new strains or ties hwater surface water used as an Hyc
ter and lower plants (whole or- sms) used to breed new strains or cties the water used as an Hyc
hwater surface water used as an Hyc
gy source
ace water for drinking Dri
ace water used as a material Indi- i-drinking purposes)
ering and attenuation of mass ement
trol of water erosion rates
Fire
ulation of peak flows Floo
ulation runoff and base flows Flov

Riparian ecosystems	Carbon sequestration by vegetation	Evaporative cooling by trees	Pollination	Providing habitats	Habitats for native pest control agents	Removal of nutrients in runoff		Filtration and storage of particles, nutrient removal	Areas of natural beauty	Sites of specific scientific interest, sites used for conservation	Heritage, sites of cultural importance	Artistic productions	Cont.
Lakes	Climate Regulation					Water purification		Nutrient and pollutant retention by macrophyte beds					
Rivers		Local climate regulation		Maintenance of nursery pop- ulations		Water purification	Dilution of wastewater	Water purification	Areas of natural beauty, ornamental use of freshwater pearls, fishes	Science and education		Artistic productions	
Class	Regulation of chemical composition of atmosphere and oceans, including maintaining rainfall patterns through evapotranspiration at the sub-continen- tal scale	Regulation of temperature and humidity, including ventilation and transpiration at local scales	Pollination	Maintaining or regulating nursery popu- lations and habitats or breeding grounds (Includes gene pool protection)	Disease control	Regulation of the chemical condition of freshwaters by living processes	Dilution or transport of wastes by fresh- water and marine ecosystems	Bio-remediation by micro-organisms, algae, plants, and animals	Elements of living systems that enable aesthetic experiences	Elements of living systems that enable scientific investigation or the creation of traditional ecological knowledge	Elements of living systems that are resonant in terms of culture or heritage	Elements of living systems used for entertainment or representation outside the setting concerned	
Group	Atmospheric compo- sition and conditions	Atmospheric compo- sition and conditions	Lifecycle mainte- nance, habitat and gene pool protection	Lifecycle mainte- nance, habitat, and gene pool protection	Pest and disease control	Water conditions	Mediation of waste, toxics and other nui- sances by non-living processes	Mediation of wastes or toxic substances of anthropogenic origin by living processes	Direct, in-situ and outdoor interactions with living systems that depend on pres-	ence in the environ- mental setting	T	Indirect, interactions with living systems	
Division	Regulation of physical, chemical, biological conditions						Transformation of biochemical or physical inputs to ecosystems		Intellectual and repre- sentative interactions with natural environ- ment				
Section									Cultural/ Non-material				

Freshwater Ecosystem Services

Cont.

Section	Division	Group	Class	Rivers	Lakes	Riparian ecosystems
	Physical and expe- riential interactions with natural environ- ment	Direct, in-situ and outdoor in- teractions with living systems that depend on presence in the environmental setting, i.e. broadly recreational activities	Elements of living systems that enable activities promoting health, recuperation or enjoyment through passive or observational inter- actions	Wildlife tourism	Wildlife tourism	Wildlife tourism and ecotourism
		Physical and experiential interac- tions with natural environment	Direct, in-situ and outdoor in- teractions with living systems that depend on presence in the environmental setting, i.e. broadly recreational activities	Angling and water sports	Angling and water sports	Sports activities
	Spiritual, symbolic and other cultural in- teractions with natural environment	Elements of living systems that are indirectly appreciated and have significance for people without their presence in the environmen- tal setting.	Elements of living systems that have spiritual or religious meaning	Religion		Totemic species or settings of religious interest
			Elements of living systems that have symbolic meaning, capture the distinctiveness of settings or their sense of place	Sense of place		Species, habitats, or landscapes that can be used as symbols
		Other biophysical characteristics of species or ecosystems that are appreciated in their own right by people	Elements or features of living systems whose contemporary exis- tence or conservation is important to people			Wilderness areas
		Other biophysical characteristics of species or ecosystems that are appreciated in their own right by people	Elements or features of living systems whose inter-generation- al existence or conservation is important to people.			Bequest

Limnetica, 45(1): 00-00 (2026)

Cont.

knowledge and that we have little information for intermediate vegetation types (Riis et al., 2020). A study on large lakes mentioned that quantification was limited to ecosystem services with commercial value due to a lack of data (Sterner et al., 2020). The scale of supply of ecosystem service often makes assessments challenging as many water ecosystem services depend on processes occurring across scales from the watershed to the habitat. Cultural services are challenging to assess in freshwater as well as other ecosystems. The multitude of notions and non-material benefits grouped into this category, which range from aesthetic to moral, leads to a focus on services with more tangible effects namely recreation and tourism (Kadykalo et al., 2019; Small et al., 2017). The role of culture in influencing the values beneficiaries attribute to ecosystem services remains poorly studied despite influencing how different groups prioritize services and manage ecosystems (Kadykalo et al., 2019; Small et al., 2017).

II. RESILIENCE OF FRESHWATER ECOSYSTEM SERVICES

Definition of resilience

The term resilience has been used in the ecological, and social sciences literature and by broader society in a variety of ways. Generally, resilience is associated with the ability to recover to a previous good state from a disturbance, trauma or pressure (Cambridge Advanced Learner's Dictionary & Thesaurus, 2024). This meaning aligns closely with the definition of engineering resilience in the ecological literature, which is which refers to the rate at which a system returns to the reference condition following a perturbation (Pimm, 1984; Van Meerbeek et al., 2021). There is an additional definition in the ecological literature which emphasises the ability to absorb disturbance and remain in the same state. Ecological resilience refers to the magnitude of disturbance that can be absorbed before the system changes its structure by changing the variables and processes that control behaviour, i.e., the ability of the system to remain in the same domain of attraction or stable state (Holling, 1973; Van Meerbeek et al., 2021).

Thus, engineering resilience generally focuses on dynamics within a stable state while ecological resilience focuses on dynamics across different stable states.

Resilience is mostly perceived as a positive aspect, however, resilience may be a positive or negative property depending on whether a given stable state is desirable in a given management context (Standish et al., 2014). Ecosystems degraded by anthropogenic activities may be as resilient or even more so than ecosystems in a better state (Durance et al., 2016; Standish et al., 2014). In these cases, resilience is a negative property, designated as unhelpful resilience, that keeps ecosystems in an undesirable state posing a management challenge (Standish et al., 2014). Regime shifts may be largely irreversible or require large amounts of energy or disturbance to shift the system to a more desirable stable state (Folke et al., 2004). The reversibility of regime shifts depends on the strength of the dominant system feedback.

Resilience mechanisms: What brings about re-silience?

Resilience can be measured at the various levels of biological organization, from population to community to ecosystem (Donohue et al., 2016; Oliver et al., 2015). However, the ability to maintain ecosystem structure and functioning and ensure the stability of ecosystem service supply is essential for ecosystem management under environmental change (Oliver et al., 2015).

The resilience of ecosystem functioning and services depends on the nature of the service, abiotic and biotic factors, and the social and governance context. This ability is shaped by the spatial and temporal scales of disturbance, and by species-to-landscape level mechanisms, including intraspecific diversity, community functional diversity and redundancy, food web complexity, and landscape heterogeneity (Gutiérrez-Cánovas et al., 2021; Oliver et al., 2015; Standish et al., 2014). Diversity, particularly functional diversity, is expected to contribute to resilience by stabilizing ecosystem functioning in the face of disturbance through compensatory dynamics or insurance effects of redundancy (de Bello et al., 2021).

Some freshwater ecosystem services directly

arise from biodiversity, at the species, taxonomic group, community or ecosystem level, such as fisheries (Brooks et al., 2016; Lynch et al., 2023). However, the relationship between biodiversity and ecosystem services is often more complex and multi-layered (Mace et al., 2012). Biodiversity may have a regulating role in key ecosystem processes, be a final ecosystem service, or a good (Mace et al., 2012). Biodiversity may be a factor controlling the ecosystem processes underpinning ecosystem services, for instance, higher biodiversity is associated with increased ecosystem functions (Mace et al., 2012). Biodiversity may also be a final service in instances where diversity contributes to goods and services, such as benefits to wild medicines and bioprospection arising from genetic diversity (Mace et al., 2012). Biodiversity itself may also be valued by humans due to its conservation or cultural value, where maintaining the diversity of wildlife, or charismatic species is considered important for recreational, educational, religious, or spiritual reasons (Mace et al., 2012).

For many ecosystem services, the contribution of biodiversity is not well defined and the evidence on biodiversity effects is mixed (Cardinale et al., 2012). The effects of biodiversity depend on the ecosystem service, whether supply or benefit is analysed, spatial scale, and on the type of linkage considered (e.g., spatial, functional, or management) (Ricketts et al., 2016). For many services, there isn't sufficient data to evaluate the relationship between biodiversity and the service (Cardinale et al., 2012). For provisioning services, service supply depends on the abundance or yield of harvested species which may or may not increase with biodiversity (Ricketts et al., 2016). For a small number of ecosystem services current evidence runs counter to general expectations, showing no relationship or negative effects of biodiversity (Cardinale et al., 2012). For instance, tree stem density, biomass, and age may have negative effects on freshwater provision through increased evapotranspiration. Nevertheless, these attributes may improve other services such as atmospheric regulation (Harrison et al. 2014).

While the evidence on the relationship between biodiversity and ecosystem services has grown it remains a critical challenge to understand how various biodiversity facets contribute the resilience of ecosystem services under environmental change.

Resilience of freshwater ecosystem services

Although an increasing number of studies reference resilience, a comprehensive understanding of ecosystem resilience, including the resilience of ecosystem services, is still lacking (Strickland et al., 2024). Few studies have measured the stability of ecosystem functions or services (Donohue et al., 2016; Oliver et al., 2015; Strickland et al., 2024). The concept of resilience is often referenced as an objective for ecosystem management or sustainable development; however, it is frequently not quantified or operational (Lloret et al., 2024). The numerous definitions of resilience applied in the realms of natural and social sciences and the diversity of indices to assess resilience even within the natural sciences have contributed to this lack of coherent global understanding (Lloret et al., 2024; Runge et al., 2025). In freshwaters, the resilience and alternative states of lake ecosystems have contributed with key references to the resilience scholarship (Carrier-Belleau et al., 2022). However, it is still unclear how widespread are regime shifts in freshwaters and challenging to find evidence of the resilience of freshwater ecosystem services.

I carried out a literature search and screening for peer-reviewed articles that studied freshwater ecosystem services resilience using Web of Science (ClarivateTM, Web of ScienceTM) to examine the current state of the literature. A search for topics "freshwater" AND "ecosystem services" AND "resilience" returned 145 results (Clarivate[™], Web of Science[™], 1 April 2025). Of these 102 articles were considered potentially relevant and the full text was analysed to understand the objectives and how they examined the resilience of freshwater ecosystem services. The articles considered irrelevant included studies focusing on marine, coastal, brackish water ecosystems, artificial water bodies, and presentations of special issues or projects. Five additional articles were not evaluated due to lack of access to the full text. Of the relevant articles almost, half were theoretical or conceptual including reviews, synthesis, perspectives, and conceptual frameworks. A few observational studies measured resilience at the level of biotic communities (Burthe et al., 2016; Feio et al., 2015). Some studies examined the resilience of local populations in relation to their dependence on ecosystem services, focusing on social contexts and governance without quantitatively measuring resilience (Komugabe-Dixson et al., 2019; Kosamu et al., 2022). One study directly connected changes in habitat for fish populations with variability in current and future spatiotemporal patterns of recreational fishery services (Cline et al., 2022). However, this example lacks metrics that connect to engineering or ecological resilience definitions. The small number of studies addressing resilience and freshwater ecosystem services, diverse methodologies and lack of operational use of resilience concepts hinder a systematic review or meta-analysis.

Therefore, the following sections present narrative examples from different freshwater ecosystems displaying diverse responses of socio-ecological ecosystems to environmental changes and disturbances. Although this a qualitative assessment, it enables a broader discussion about the resilience of freshwater socio-ecological systems and how close they are to tipping points and regime shifts.

African lakes

The great African lakes hold an estimated 25% of the world's liquid surface freshwater. African lakes contribute to the livelihoods of millions of people as well as national economies. Some lakes experience fluctuations in water levels as well as dry-out periods which impact the livelihoods of local communities (Ogutu-Ohwayo et al., 2016). The increasing exploitation of lake resources during the past century has led to habitat loss and degradation, introduction of exotic species, pollution from agricultural, industrial and urban runoff (Kafumbata et al., 2014; Ogutu-Ohwayo et al., 2016). These stressors are likely to be aggravated by climate change (Mutanda & Nhamo, 2024). The loss of ecosystem services has profound implications for the local communities which rely heavily on the flow of ecosystem services from the lakes for food security and employment.

In Lake Naivasha in Kenya population growth caused riparian vegetation degradation, high nutrient, sediment loads and increasing water abstraction for agriculture and industry (Mutethya & Yongo, 2021; Renaut & Owen, 2023). The lake levels have decreased because of the increasing water demand for a growing horticultural industry and population and it has become eutrophic due to nutrient inputs (Renaut & Owen, 2023). The introduction of non-native fish may have contributed further to alteration of the lake's trophic state (Mutethya & Yongo, 2021). The lake levels have recovered following drought and its ecosystem services have remained resilient due to groundwater recharge from aquifers and the establishment of commercial fisheries around non-native fish species (Harper et al., 2011; Kafumbata et al., 2014; Mutethya & Yongo, 2021).

Lake Chilwa in Malawi on the other hand is showing signs of declining resilience (Kafumbata et al., 2014). The lake experiences significant inter- and intra-annual variations in water levels including severe lake level recessions and complete drying associated with droughts (Kambombe et al., 2021). The lake recovered from desiccation; however, it has a negative water budget and the wetland area is decreasing due to deforestation and agricultural expansion (Kambombe et al., 2023; Njava et al., 2011). Fisheries are strongly dependent on lake levels, declining in periods of lake recession and requiring three to four years to recover after a severe recession (Rebelo et al., 2011; Njaya et al., 2011). Since fishing may contribute up to half of income for local populations, lake recessions have a significant impact on local livelihoods (Njaya et al., 2011). During dry periods, communities turn to other activities leading to increased exploitation of exposed land for agriculture, livestock grazing, and wetland birds (Kafumbata et al., 2014; Njaya et al., 2011). The loss of income leads to migration of fishermen while women and children are left with minimal resources in a context of increasing conflicts for the exploitation of resources (Nagoli & Chiwona-Karltun, 2017).

Lake Chad in West Africa, in the conjunction of Chad, Cameroon, Nigeria and Niger, is considered to have shifted to an alternative ecosystem state since this once-large lake has shrunk by more than 90% since the 1960s splitting into two pools (Bouchez et al., 2016; Leblanc et al., 2011). The dramatic change in the lake is attributed to declines in precipitation and streamflow associated with climate change as well as water abstraction for agriculture in the tributaries (Pham-Duc et al., 2020). The lake is a source of freshwater and natural resources for two million people on the lakeshore and over ten million further afield (Lake Chad Basin Commission, 2016; Riebe & Dressel, 2021). The decline of lake levels reduced ecosystem services supply, namely water supply, crop production, livestock production, and fisheries (Okpara et al., 2016; Riebe & Dressel, 2021). This has led to food insecurity and loss of income for local communities which migrate within the lake and change their activities, often leading to conflicts due to competition for limited resources (Okpara et al., 2016). The instability and unemployment caused by the declines of Lake Chad are believed to have contributed to the increasing the influence of the armed group Boko Haram in the region (Owonikoko & Momodu, 2020).

The three African lakes presented here provide three examples of various levels of resilience of ecosystem services. Lake Naivasha retained resilience thanks to groundwater recharge, while Lake Chilwa is in a downward spiral with declining capacity to provide ecosystem services (Kafumbata et al., 2014). Lake Chad is considered to have passed a critical threshold into another state that cannot provide adequate ecosystem services to communities, contributing to large-scale conflicts in the region (Bouchez et al., 2016; Kafumbata et al., 2014). In all three examples, local communities are strongly dependent on the ecosystem services provided by the lakes and thus food security and livelihoods are highly sensitive to variability and shifts in the natural systems. This is exacerbated by the lack of adequate human responses, particularly institutional responses that mitigate impacts on local communities (Kafumbata et al., 2014).

The Pantanal Wetland in Brazil

The Pantanal is the world's largest freshwater wetland and a UNESCO World Heritage Site occupying an area of 140 000 km² mostly in Brazil (90% [Pott & Pott, 2004; UNESCO, 2025]). The Pantanal is a seasonal floodplain in the upper Paraguay River basin alternating between flooded and dry phases which influence the ecosystem functioning, biodiversity and local communities' lifestyles (Pott & Pott, 2004). The landscape is a mosaic of flooded and non-flooded grasslands, forests, savannahs, and permanent and temporary water bodies hosting high species diversity (Pott & Pott, 2004). The main source of income is traditional extensive cattle grazing, where local communities move the cattle along flooding gradients to graze using fire to manage vegetation and promote the growth of native grasses (Tomas et al., 2024). Fisheries are the second most important activity as riverside communities move through the canals and lakes to find fishing grounds (Tomas et al., 2024). Agriculture, hydroelectric developments and climate change pose major threats to the Pantanal ecosystem (Thielen et al., 2020; Wantzen et al., 2024). The Pantanal is prone to wildfires during the dry season and fires started by human activities get out of control often (Pott & Pott, 2004). In 2020 the Pantanal experienced the worst drought in 60 years leading to massive fires that burned nearly four million hectares (Ferreira Barbosa et al., 2022; Laboratório de Aplicações de Satélites Ambientais, 2021; Libonati et al., 2020). In 2024 drought conditions fuelled large fires again burning more than 2 million hectares (Laboratório de Aplicações de Satélites Ambientais, 2021). Recent studies estimate that at least 16.952 million vertebrates were killed by the 2020 fires (Tomas et al., 2021). Recurrent fire may eliminate fire-sensitive species increasing the dominance of fire-prone species (Libonati et al., 2020; Pott & Pott, 2004). However, the full-scale impacts of the extreme events of 2020 and 2024 remain largely unknown requiring additional monitoring to understand Pantanal's resilience (Tomas et al., 2021). The large fires raised awareness about the need to implement integrated fire management policies prompting the approval of state legislation to regulate fire use (Tomas et al., 2021). However, severe fire legislation that criminalises fire use by traditional communities may risk local ecosystems since cattle ranchers and farmers turn to exotic grass species (Garcia et al., 2021; Tomas et al., 2019). Overall,

we still require further research to understand the socio-ecological resilience of the Pantanal especially in the long-term.

Riparian forests of the Murray-Darling Basin in Australia

The Murray-Darling Basin is the largest freshwater basin in Australia spanning one million square kilometres of south-eastern Australia across the states of New South Wales, Queensland, South Australia, Victoria and the Australian Capital Territory (Murray-Darling Basin Authority, 2023b). The basin is responsible for 40% of Australia's agricultural production and it is home to more than two million people (Murray-Darling Basin Authority, 2023a). The ecosystems are valued nationally and internationally supporting tourism and recreation and hold cultural value for indigenous populations (Murray-Darling Basin Authority, 2023a). This region experienced a multi-year drought between 2000-2009 known as the Millenium Drought (van Dijk et al., 2013). The Millenium Drought was the most severe on record for SE Australia and, together with high water demand, led to a water availability of less than 40% of the historical average (van Dijk et al., 2013). The river ecosystems and the agriculture in the Murray-Darling Basin were severely affected by this drought, and people faced higher electricity prices and water use restrictions (van Dijk et al., 2013). The drought and water abstraction caused the dieback of iconic floodplain species along the river (Harris et al., 2018). The river red gum forests (Eucalyptus camaldulensis) showed reductions in canopy cover and mortality, particularly in higher-density forest areas (Harris et al., 2018). The recovery of these forests is hindered by declines in recruitment due to lower flooding frequency associated to river regulation, increasing salinity of soil and water (Mac Nally et al., 2011).

The drought and tree dieback caused severe declines in freshwater ecosystem services. Aesthetic values, pollination, timber, carbon storage, nutrient cycling, erosion and water quality regulation all suffered significant reductions (van Dijk et al., 2013). The losses in water ecosystem services caused by the Millennium Drought required an estimated expenditure of 810 million AUD to mitigate ecosystem services losses and to implement adaptation measures (Banerjee et al., 2013). Riverbanks receded, slumped and in some cases collapsed as the water decreased leading to the need to develop a hazard mitigation plan (Banerjee et al., 2013). Seven years after the drought one-third of the watersheds in the state of Victoria had not fully recovered to pre-drought runoff levels (Peterson et al., 2021). These trends suggest that the socio-ecological system has passed a tipping point, from a social and political perspective if not from an ecological perspective.

III. RESILIENCE OF FRESHWATER ECOSYSTEMS AND SERVICES UNDER GLOBAL CHANGE

According to the latest assessment of planetary boundaries, the freshwater boundary has already been transgressed (Richardson et al., 2023). This evaluation is supported by an increase in land area where blue (streamflow) and green water (soil moisture) levels deviate from pre-industrial reference conditions (Porkka et al., 2024). Another concerning indicator is the increase in freshwater blooms in lakes since the 1980s (Ho et al., 2019). Broadly the trajectories of freshwater ecosystems and their services under climate change are expected to depend on whether ecosystems are energy or water-limited systems (Campbell et al., 2022). In energy-limited systems in higher latitudes and mountains the increasing temperature promotes range shifts, increasing competition and possibly excluding cold-adapted species (Nilsson et al., 2012; Perry et al., 2012). Aquatic and riparian cold-water animal species are already experiencing population declines (Durance & Ormerod, 2007; Rogers et al., 2020). In water-limited systems, such as those in the Mediterranean climate, increasing water limitation will promote environmental filtering, declines in functional diversity and shifts toward more conservative strategies (Portela, Durance, et al., 2023a). Obligate riparian and drought-sensitive plant species are expected to decline and annual species to increase in semi-arid rivers (Perry et al., 2012; Stromberg et al., 2012). These trends may represent significant departures from the current ecosystem status and functional diversity levels. The ecosystem services expected to be most affected include water provisioning, fisheries, and many regulation services such as climate change mitigation and water quality regulation (Campbell et al., 2022; Portela, Durance, et al., 2023b). Since functional diversity underpins ecosystem functioning and stability we also expect changes from current baselines of ecosystem resilience (Biggs et al., 2020).

The examples discussed in this work illustrate declines in freshwater ecosystem services, erosion of resilience in socio-ecological systems as well as the uncertainty surrounding the identification of regime shifts and future trajectories. Clear examples of loss of resilience and regime shifts are still rare and mostly associated with lake systems, as exemplified here by Lake Chad. A study of LTER sites in the US over 40 years finds limited evidence of lasting regime shifts in freshwater and forest ecosystems (Campbell et al., 2022). This may be because many of these LTER sites are still experiencing conditions similar to their historical climate space (Campbell et al., 2022). It is also hypothesised that shallow lakes are more prone to regime shifts whereas running waters are more resilient (Durance et al., 2016). However, the observational or experimental evidence available on the resilience of freshwater ecosystems and their services remains limited. Many studies discuss resilience and associated topics in the introduction or discussion, but do not aim to quantify it (Carrier-Belleau et al., 2022). This prevents a comprehensive, systematic assessment of the resilience of freshwater ecosystems and their services globally or comparatively. While the freshwater resilience research continues to evolve, there are examples of warning signs of loss of resilience particularly in response to extreme climate events and disturbances.

Disturbances such as drought, fire, hurricanes, and ice storms appear to be pushing socio-ecological systems near or beyond tipping points and towards new stable states. Disturbances across land and water can impact and destabilise freshwater ecosystems and their services (Crausbay et al., 2020; Falkenmark & Wang-Erlandsson, 2021). The Millenium Drought in Australia, the Paranal fires, and the recurrent drying in Lake Chilwa are examples of recurring or prolonged extremes that may be overwhelming the capacity of socio-ecological systems to recover and sustain ecosystem services. Furthermore, extreme events can cause legacy effects in key attributes of ecosystem functioning, where declines in ecosystem productivity and streamflow last several years following the event (Peterson et al., 2021; Portela, Goncalves, et al., 2023). However, their effects may not be fully evident at short time scales. A watershed scale experiment at a LTER site suggests initial recoveries of ecosystem functions following disturbance do not preclude later regime shifts as other disturbances impact recovery trajectories (Jackson et al., 2018). Climate extremes and other disturbances may be as important or even more so than changes in average climate for the resilience of freshwater ecosystems in the future.

The resilience of the socio-ecological systems is also modulated by social systems including community and institutional dynamics as shown by the examples presented here. The Millennium Drought illustrates how high water consumption and river regulation contribute to the decline of riparian forests and their limited recovery. The examples from African lakes and the Australian Millennium Drought offer a stark contrast on institutional dynamics. In Lake Chad the lack of institutional responses has contributed to unemployment, conflicts, and instability in the region as the livelihoods of the people strongly depend on ecosystem services provided by or connected to the lake. In the Australian Drought the stronger institutional capacity allowed the implementation of several measures to mitigate and adapt to the effects of the event, even if at a significant cost. The loss of water-resilience has been linked with broader social dynamics including armed conflicts, uprisings, and even the collapse of ancient civilizations (Falkenmark & Wang-Erlandsson, 2021). These connections are still debated and difficult to ascertain due to the complexity of social systems. Nevertheless, the communities most affected by loss of freshwater ecosystem services are undoubtedly those that depend the most on ecosystems. Then, social structures at individual, community and institutional levels shape the capacity to adapt or replace ecosystem services with technological solutions or grey infrastructure thus mediating the resilience of freshwater social-ecological systems (Falkenmark et al., 2019).

The resilience of freshwaters to climate change, compounding disturbances and stressors is still unclear and emerging, thus further research is needed the resilience of ecosystems and their services. First, we need to address the basic gaps in the assessment of freshwater ecosystem services and their linkages with biodiversity. Second, we need to improve our understanding of resilience, identifying alternative stable states and tipping points in freshwaters, especially in running waters. To that end, we need long-term studies and monitoring as well as retrospective studies to understand the response of ecosystems to past and ongoing changes and identify potential tipping points. Third, we need to enhance our understanding of the roles of social dynamics in modulating resilience and to include multidisciplinary perspectives in evaluating freshwater ecosystem services and resilience.

CONCLUSION

In conclusion, this work illustrates diverse responses of freshwater socio-ecological systems to environmental changes, highlighting warning signs of declining resilience due to climate change and altered disturbance regimes. The examples highlight how significant the consequences for social systems can be as well as the role of social structures in mediating resilience. However, a high degree of uncertainty still surrounds the identification of tipping points, regime shifts, and future ecosystem trajectories. The responses of freshwater ecosystems to climate change, multiple disturbances and stressors are still emerging. Predicting the dynamics of these critical socio-ecological systems in the future requires long-term studies directly examining resilience. Research on the resilience of freshwaters is essential to ensure they continue providing critical ecosystem services to society, thereby fostering resilient communities.

ACKNOWLEDGEMENTS

This work was supported by Fundação para a Ciência e Tecnologia through the Doctoral fellowship SFRH/BD/115030/2016 funded by Fundação para a Ciência e Tecnologia, the Ministério da Ciência, Tecnologia e Ensino Superior, the European Social Fund, Human Capital Operational Program (POCH), and North Portugal Regional Operational Programme (NORTE2020). The article was developed in the context of the XV Limnology Research Award for best doctoral thesis defended in 2023 by the Iberian Association of Limnology.

REFERENCES

- Aylward, B., Börkey, P., & Allan, T. (2005). Freshwater Ecosystem Services. In K. Chopra, R. Leemans, P. Kumar, & H. Simons (Eds.), Ecosystems and human well-being : policy responses : findings of the Responses Working Group of the Millennium Ecosystem Assessment (Vol. 3).
- Banerjee, O., Bark, R., Connor, J., & Crossman, N. D. (2013). An ecosystem services approach to estimating economic losses associated with drought. *Ecological Economics*, 91, 19-27. DOI: 10.1016/j.ecolecon.2013.03.022
- Biggs, C. R., Yeager, L. A., Bolser, D. G., Bonsell, C., Dichiera, A. M., Hou, Z., Keyser, S. R., Khursigara, A. J., Lu, K., Muth, A. F., Negrete, B., & Erisman, B. E. (2020). Does functional redundancy affect ecological stability and resilience? A review and meta-analysis. *Ecosphere*, 11(7). DOI: 10.1002/ecs2.3184
- Bouchez, C., Goncalves, J., Deschamps, P., Vallet-Coulomb, C., Hamelin, B., Doumnang, J. C., & Sylvestre, F. (2016). Hydrological, chemical, and isotopic budgets of Lake Chad: a quantitative assessment of evaporation, transpiration and infiltration fluxes. *Hydrology and Earth System Sciences*, 20(4), 1599-1619. DOI: 10.5194/hess-20-1599-2016
- Braat, L. C., & de Groot, R. (2012). The ecosystem services agenda:bridging the worlds of natural science and economics, conservation and development, and public and private policy. *Ecosystem Services*, 1(1), 4-15. DOI: 10.1016/j.ecoser.2012.07.011
- Brauman, K. A. (2015). Hydrologic ecosystem services: linking ecohydrologic processes to human well-being in water research and watershed management. *Wiley Interdisciplinary Reviews-Water*, 2(4), 345-358. DOI: 10.1002/

wat2.1081

- Brauman, K. A., Garibaldi, L. A., Polasky, S., Aumeeruddy-Thomas, Y., Brancalion, P. H. S., DeClerck, F.,Verma, M. (2020). Global trends in nature's contributions to people. *Proceedings of the National Academy of Sciences*, 117(51), 32799-32805. DOI: 10.1073/ pnas.2010473117
- Brooks, E. G. E., Holland, R. A., Darwall, W. R. T., & Eigenbrod, F. (2016). Global evidence of positive impacts of freshwater biodiversity on fishery yields. *Global Ecology and Biogeography*. DOI: 10.1111/geb.12435
- Burthe, S. J., Henrys, P. A., Mackay, E. B., Spears, B. M., Campbell, R., Carvalho, L., Dudley, B.,Daunt, F. (2016). Do early warning indicators consistently predict nonlinear change in long-term ecological data? *Journal of Applied Ecology*, 53(3), 666-676. DOI: 10.1111/1365-2664.12519
- Cambridge Advanced Learner's Dictionary & Thesaurus. (2024). *Definition of resilience*. https://dictionary.cambridge.org/dictionary/ english/resilience
- Campbell, J. L., Driscoll, C. T., Jones, J. A., Boose, E. R., Dugan, H. A., Groffman, P. M., Jackson, C. R.,Zimmerman, J. K. (2022). Forest and Freshwater Ecosystem Responses to Climate Change and Variability at US LTER Sites. *Bioscience*, 72(9), 851-870. DOI: 10.1093/biosci/biab124
- Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., Naeem, S. (2012). Biodiversity loss and its impact on humanity. *Nature*, 486(7401), 59-67. DOI: 10.1038/nature11148
- Carrier-Belleau, C., Pascal, L., Nozais, C., & Archambault, P. (2022). Tipping points and multiple drivers in changing aquatic ecosystems: A review of experimental studies [Review]. *Limnology and Oceanography*, 67, S312-S330. DOI: 10.1002/lno.11978
- Cline, T. J., Muhlfeld, C. C., Kovach, R., Al-Chokhachy, R., Schmetterling, D., Whited, D., & Lynch, A. J. (2022). Socioeconomic resilience to climatic extremes in a freshwater fishery. *Science Advances*, 8(36), eabn1396. DOI: 10.1126/sciadv.abn1396

Crausbay, S. D., Betancourt, J., Bradford, J.,

.....Carter, S. (2020). Unfamiliar Territory: Emerging Themes for Ecological Drought Research and Management. *One Earth*, 3(3), 337-353. DOI: 10.1016/j.oneear.2020.08.019
Datry, T., Boulton, A. J., Bonada, N., Fritz, K., Leigh, C., Sauquet, E., Tockner, K., Hugueny,

Leigh, C., Sauquet, E., Tockner, K., Hugueny, B., & Dahm, C. N. (2018). Flow intermittence and ecosystem services in rivers of the Anthropocene. *Journal of Applied Ecology*, 55(1), 353-364. DOI: 10.1111/1365-2664.12941

Cartwright, J., Dennison, W. C., Dunham, J.,

- de Bello, F., Lavorel, S., Hallett, L. M., Valencia, E., Garnier, E., Roscher,Leps, J. (2021). Functional trait effects on ecosystem stability: assembling the jigsaw puzzle. *Trends in Ecology & Evolution*, 36(9), 822-836. DOI: 10.1016/j.tree.2021.05.001
- Díaz, S., Demissew, S., Carabias, J., Joly, C., Lonsdale, ...Zlatanova, D. (2015). The IPBES Conceptual Framework — connecting nature and people. *Current Opinion in Environmental Sustainability*, 14, 1-16. DOI: 10.1016/j. cosust.2014.11.002
- Díaz, S., Pascual, U., Stenseke, M., Martín-López, B., Watson, R. T., ...Shirayama, Y. (2018). Assessing nature's contributions to people. *Science*, 359(6373), 270-272. DOI: 10.1126/science.aap8826
- Donohue, I., Hillebrand, H., Montoya, J. M., Petchey, O. L., Pimm, S. L., Fowler, M. S.,Yang, Q. (2016). Navigating the complexity of ecological stability. *Ecology Letters*, 19(9), 1172-1185. DOI: 10.1111/ele.12648
- Dunham, J. B., Angermeier, P. L., Crausbay, S. D., Cravens, A. E., Gosnell, H., McEvoy, J., Moritz, M. A., Raheem, N., & Sanford, T. (2018). Rivers are social–ecological systems: Time to integrate human dimensions into riverscape ecology and management. *WIRES Water*, 5(4). DOI: 10.1002/wat2.1291
- Durance, I., Bruford, M. W., Chalmers, R., Chappell, N. A., Christie, M., Cosby, B.
 J., Noble, D., Ormerod, S. J., Prosser, H., Weightman, A., & Woodward, G. (2016). The Challenges of Linking Ecosystem Services to Biodiversity. *Academic Press*, 54, 87-134. DOI: 10.1016/bs.aecr.2015.10.003 (*Advances in Ecological Research*)
- Durance, I., & Ormerod, S. J. (2007). Climate

change effects on upland stream macroinvertebrates over a 25-year period. *Global Change Biology*, 13(5), 942-957. DOI: 10.1111/j.1365-2486.2007.01340.x

- Egoh, B., Drakou, E. G., Dunbar, M. B., Maes, J., & Willemen, L. (2012). Indicators for mapping ecosystem services: a review. *JRC Publications Repository*. DOI: 10.2788/41823
- Ehrlich, P., & Ehrlich, A. (1981). *Extinction: the causes and consequences of the disappearance of species*. Random House. New York, 72-98
- Falkenmark, M., & Wang-Erlandsson, L. (2021). A water-function-based framework for understanding and governing water resilience in the Anthropocene [Review]. One Earth, 4(2), 213-225. DOI: 10.1016/j.oneear.2021.01.009
- Falkenmark, M., Wang-Erlandsson, L., & Rockström, J. (2019). Understanding of water resilience in the Anthropocene. *Journal of Hydrology X*, 2, 100009. DOI: 10.1016/j.hydroa.2018.100009
- Feio, M. J., Dolédec, S., & Graça, M. A. S. (2015). Human disturbance affects the long-term spatial synchrony of freshwater invertebrate communities. *Environmental Pollution*, 196, 300-308. DOI: 10.1016/j.envpol.2014.09.026
- Ferreira Barbosa, M. L., Haddad, I., da Silva Nascimento, A. L., Máximo da Silva, G., Moura da Veiga, R., Hoffmann, T. B., Oighenstein Anderson, L. (2022). Compound impact of land use and extreme climate on the 2020 fire record of the Brazilian Pantanal. *Global Ecology and Biogeography*, 31(10), 1960-1975. DOI: 10.1111/geb.13563
- Ferreira, V., Albariño, R., Larrañaga, A., LeRoy, C. J., Masese, F. O., & Moretti, M. S. (2022). Ecosystem services provided by small streams: an overview. *Hydrobiologia*, 850(12-13), 2501-2535. DOI: 10.1007/s10750-022-05095-1
- Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L., & Holling, C. S. (2004). Regime Shifts, Resilience, and Biodiversity in Ecosystem Management. Annual *Review of Ecology, Evolution, and Systematics*, 35(1), 557-581. DOI: 10.1146/annurev. ecolsys.35.021103.105711

Forzieri, G., Dakos, V., McDowell, N. G.,

Ramdane, A., & Cescatti, A. (2022). Emerging signals of declining forest resilience under climate change. *Nature*, 608(7923), 534-539. DOI: 10.1038/s41586-022-04959-9

- Garcia, L. C., Szabo, J. K., de Oliveira Roque, F., de Matos Martins Pereira, A., Nunes da Cunha, C., Damasceno-Junior, G. A., Morato, R. G., Tomas, W. M., Libonati, R., & Ribeiro, D. B. (2021). Record-breaking wildfires in the world's largest continuous tropical wetland: Integrative fire management is urgently needed for both biodiversity and humans. *Journal* of Enviromental Management, 293, 112870. DOI: 10.1016/j.jenvman.2021.112870
- Gómez-Baggethun, E., de Groot, R., Lomas, P. L., & Montes, C. (2010). The history of ecosystem services in economic theory and practice: From early notions to markets and payment schemes. *Ecological Economics*, 69(6), 1209-1218. DOI: 10.1016/j.ecolecon.2009.11.007
- Greenwood, E. E., Lauber, T., van den Hoogen, J., Donmez, A., Bain, R. E. S., Johnston, R., Crowther, T. W., & Julian, T. R. (2024). Mapping safe drinking water use in low- and middle-income countries. *Science*, 385(6710), 784-790. DOI: 10.1126/science.adh9578
- Grizzetti, B., Lanzanova, D., Liquete, C., Reynaud, A., & Cardoso, A. C. (2016). Assessing water ecosystem services for water resource management. *Environmental Science* & *Policy*, 61, 194-203. DOI: 10.1016/j. envsci.2016.04.008
- Grizzetti, B., Liquete, C., Pistocchi, A., Vigiak, O., Zulian, G., Bouraoui, F., De Roo, A., & Cardoso, A. C. (2019). Relationship between ecological condition and ecosystem services in European rivers, lakes and coastal waters. *Science of the Total Environment*, 671, 452-465. DOI: 10.1016/j.scitotenv.2019.03.155
- Gutiérrez-Cánovas, C., Worthington, T. A., Jâms,
 I. B., Noble, D. G., Perkins, D. M., Vaughan, I.
 P., Woodward, G., Ormerod, S. J., & Durance,
 I. (2021). Populations of high-value predators reflect the traits of their prey. *Ecography*, 44(5), 690-702. DOI: 10.1111/ecog.05438
- Haines-Young, R. (2023). Common International Classification of Ecosystem Services (CICES) V5.2 and Guidance on the Application of the Revised Structure. www.cices.eu

- Hanna, D. E. L., Tomscha, S. A., Ouellet Dallaire, C., Bennett, E. M., & Hooftman, D. (2018). A review of riverine ecosystem service quantification: Research gaps and recommendations. *Journal of Applied Ecology*, 55(3), 1299-1311. DOI: 10.1111/1365-2664.13045
- Harper, D. M., Morrison, E. H. J., Macharia, M. M., Mavuti, K. M., & Upton, C. (2011). Lake Naivasha, Kenya: Ecology, Society and Future. *Freshwater Reviews*, 4(2), 89-114, 126. DOI: 10.1608/FRJ-4.2.149
- Harris, R. M. B., Beaumont, L. J., Vance, T. R., Tozer, C. R., Remenyi, T. A., Perkins-Kirkpatrick, S. E.,Bowman, D. M. J. S. (2018). Biological responses to the press and pulse of climate trends and extreme events. *Nature Climate Change*, 8(7), 579-587. DOI: 10.1038/s41558-018-0187-9
- Heino, J., Alahuhta, J., Bini, L. M., Cai, Y., Heiskanen, A. S., Hellsten, S., Kortelainen, P., Kotamaki, N., Tolonen, K. T., Vihervaara, P., Vilmi, A., & Angeler, D. G. (2021). Lakes in the era of global change: moving beyond single-lake thinking in maintaining biodiversity and ecosystem services. Biological Reviews Cambridge Philosophical Society, 96(1), 89-106. DOI: 10.1111/brv.12647
- Ho, J. C., Michalak, A. M., & Pahlevan, N. (2019). Widespread global increase in intense lake phytoplankton blooms since the 1980s. *Nature*, 574(7780), 667-670. DOI: 10.1038/ s41586-019-1648-7
- Holling, C. S. (1973). Resilience and Stability of Ecological Systems. *Annual Review of Ecology and Systematics*, 4(1), 1-23. DOI: 10.1146/ annurev.es.04.110173.000245
- Holmlund, C. M., & Hammer, M. (1999). Ecosystem services generated by fish populations. *Ecological Economics*, 29(2), 253-268. DOI: 10.1016/S0921-8009(99)00015-4
- Jackson, C. R., Webster, J. R., Knoepp, J. D., Elliott, K. J., Emanuel, R. E., Caldwell, P. V., & Miniat, C. F. (2018). Unexpected ecological advances made possible by long-term data: A Coweeta example. *WIRES Water*, 5(2). DOI: 10.1002/wat2.1273
- Kadykalo, A. N., López-Rodriguez, M. D., Ainscough, J., Droste, N., Ryu, H., Ávila-Flores, G., Le Clec'h, S., Muñoz,

M. C., Nilsson, L., Rana, S., Sarkar, P., Sevecke, K. J., & Harmáčková, Z. V. (2019). Disentangling 'ecosystem services' and 'nature's contributions to people'. *Ecosystems and People*, 15(1), 269-287. DOI: 10.1080/26395916.2019.1669713

- Kafumbata, D., Jamu, D., & Chiotha, S. (2014). Riparian ecosystem resilience and livelihood strategies under test: lessons from Lake Chilwa in Malawi and other lakes in Africa [Review]. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 369(1639), 8, Article 20130052. DOI: 10.1098/rstb.2013.0052
- Kambombe, O., Ngongondo, C., Eneya, L., Monjerezi, M., & Boyce, C. (2021). Spatiotemporal analysis of droughts in the Lake Chilwa Basin, Malawi. *Theoretical and Applied Climatology*, 144(3), 1219-1231. DOI: 10.1007/s00704-021-03586-0
- Kambombe, O., Ngongondo, C., Monjerezi, M., & Eneya, L. (2023). Investigating human-induced threat to hydrological regime of Lake Chilwa Basin, Malawi. *Applied Water Science*, 13(7), 161. DOI: 10.1007/s13201-023-01965-8
- Komugabe-Dixson, A. F., de Ville, N. S. E., Trundle, A., & McEvoy, D. (2019). Environmental change, urbanisation, and socio-ecological resilience in the Pacific: Community narratives from Port Vila, Vanuatu. *Ecosystem Services*, 39, 100973. DOI: 10.1016/j.ecoser.2019.100973
- Kosamu, I. B. M., Makwinja, R., Kaonga, C. C., Mengistou, S., Kaunda, E., Alamirew, T., & Njaya, F. (2022). Application of DPSIR and Tobit Models in Assessing Freshwater Ecosystems: The Case of Lake Malombe, Malawi. *Water*, 14(4). DOI: 10.3390/w14040619
- Laboratório de Aplicações de Satélites Ambientais. (2021). *ALARMES* - Alerta de Área Queimada com Monitoramento Estimado por Satélite. https://alarmes.lasa.ufrj.br/platform/ dashboard/9091
- Lake Chad Basin Commission. (2016). *The Lake chad development and climate resilience action plan.*
- Leblanc, M., Lemoalle, J., Bader, J. C., Tweed, S., & Mofor, L. (2011). Thermal remote sensing of water under flooded vegetation: New obser-

vations of inundation patterns for the 'Small' Lake Chad. *Journal of Hydrology*, 404(1-2), 87-98. DOI: 10.1016/j.jhydrol.2011.04.023

- Libonati, R., DaCamara, C. C., Peres, L. F., Sander de Carvalho, L. A., & Garcia, L. C. (2020). Rescue Brazil's burning Pantanal wetlands. *Nature*, 588(7837), 217-219. DOI: 10.1038/d41586-020-03464-1
- Lloret, F., Hurtado, P., Espelta, J. M., Jaime, L., Nikinmaa, L., Lindner, M., & Martínez-Vilalta, J. (2024). ORF, an operational framework to measure resilience in social– ecological systems: the forest case study. *Sustainability Science*, 19(5), 1579-1593. DOI: 10.1007/s11625-024-01518-1
- Lynch, A. J., Cooke, S. J., Arthington, A. H., Baigun, C., Bossenbroek, L., Dickens, C., ... Jähnig, S. C. (2023). People need freshwater biodiversity. *WIRES Water*, 10(3). DOI: 10.1002/wat2.1633
- Mac Nally, R., Cunningham, S. C., Baker, P. J., Horner, G. J., & Thomson, J. R. (2011). Dynamics of Murray-Darling floodplain forests under multiple stressors: The past, present, and future of an Australian icon. *Water Resources Research*, 47(12). DOI: 10.1029/2011wr010383
- Mace, G. M., Norris, K., & Fitter, A. H. (2012). Biodiversity and ecosystem services: a multilayered relationship. *Trends in Ecology & Evolution*, 27(1), 19-26. DOI: 10.1016/j. tree.2011.08.006
- Maes, J., Egoh, B., Willemen, L., Liquete, C., Vihervaara, P., Schägner, J. P., Bidoglio, G. (2012). Mapping ecosystem services for policy support and decision making in the European Union. *Ecosystem Services*, 1(1), 31-39. DOI: 10.1016/j.ecoser.2012.06.004
- Maltby, E., Ormerod, S., Acreman, M., Dunbar, M., Jenkins, A., Maberly, S., Newman, J., Blackwell, M., & Ward, R. (2011). Freshwaters : openwaters, wetlands and floodplains [chapter 9]. In UK National Ecosystem Assessment: understanding nature's value to society. Technical Report (pp. 295-360). UNEP-WC-MC. http://nora.nerc.ac.uk/16133/
- Millennium Ecosystem Assessment. (2005). *Eco*systems and Human Well-being: Synthesis. Island Press.

- Murray-Darling Basin Authority. (2023a, 11 December 2023). *The Basin*. https://www.mdba. gov.au/basin
- Murray-Darling Basin Authority. (2023b, 29 November 2023). *Basin Plan*. https://www.mdba. gov.au/water-management/basin-plan
- Mutanda, G. W., & Nhamo, G. (2024). Impact of climate change on Africa's major lakes: a systematic review incorporating pathways of enhancing climate resilience. *Frontiers in Water*, 6. DOI: 10.3389/frwa.2024.1443989
- Mutethya, E., & Yongo, E. (2021). A comprehensive review of invasion and ecological impacts of introduced common carp (Cyprinus carpio) in Lake Naivasha, Kenya. *Lakes & Reservoirs: Science, Policy and Management* for Sustainable Use, 26(4), e12386. DOI: 10.1111/lre.12386
- Nagoli, J., & Chiwona-Karltun, L. (2017). Uncovering human social networks in coping with Lake Chilwa recessions in Malawi. *Journal of Environmental Management*, 192, 134-141. DOI: 10.1016/j.jenvman.2016.12.049
- Nicolás Ruiz, N., Suárez Alonso, M. L., & Vidal-Abarca, M. R. (2021). Contributions of dry rivers to human well-being: A global review for future research. *Ecosystem Services*, 50, 101307. DOI: 10.1016/j.ecoser.2021.101307
- Nilsson, C., Jansson, R., Kuglerová, L., Lind, L., & Ström, L. (2012). Boreal Riparian Vegetation Under Climate Change. *Ecosystems*, 16(3), 401-410. DOI: 10.1007/s10021-012-9622-3
- Njaya, F., Snyder, K. A., Jamu, D., Wilson, J., Howard-Williams, C., Allison, E. H., & Andrew, N. L. (2011). The natural history and fisheries ecology of Lake Chilwa, southern Malawi. *Journal of Great Lakes Research*, 37, 15-25. DOI: 10.1016/j.jglr.2010.09.008
- Ogutu-Ohwayo, R., Natugonza, V., Musinguzi, L., Olokotum, M., & Naigaga, S. (2016). Implications of climate variability and change for African lake ecosystems, fisheries productivity, and livelihoods. *Journal of Great Lakes Research*, 42(3), 498-510. DOI: 10.1016/j. jglr.2016.03.004
- Okpara, U. T., Stringer, L. C., & Dougill, A. J. (2016). Lake drying and livelihood dynamics

in Lake Chad: Unravelling the mechanisms, contexts and responses. *Ambio*, 45(7), 781-795. DOI: 10.1007/s13280-016-0805-6

- Oliver, T. H., Heard, M. S., Isaac, N. J. B., Roy, D. B., Procter, D., Eigenbrod, F., & Bullock, J. M. (2015). Biodiversity and Resilience of Ecosystem Functions. Trends in Ecology & Evolution, 30(11), 673-684. DOI: 10.1016/j. tree.2015.08.009
- Owonikoko, S. B., & Momodu, J. A. (2020). Environmental degradation, livelihood, and the stability of Chad Basin Region. *Small Wars & Insurgencies*, 31(6), 1295-1322. DOI: 10.1080/09592318.2020.1776092
- Pastor, A. V., Tzoraki, O., Bruno, D., Kaletová, T., Mendoza-Lera, C., Alamanos, A., Jorda-Capdevila, D. (2022). Rethinking ecosystem service indicators for their application to intermittent rivers. *Ecological Indicators*, 137. DOI: 10.1016/j.ecolind.2022.108693
- Perry, L. G., Andersen, D. C., Reynolds, L. V., Nelson, S. M., & Shafroth, P. B. (2012). Vulnerability of riparian ecosystems to elevated CO2 and climate change in arid and semiarid western North America. *Global Change Biology*, 18(3), 821-842. DOI: 10.1111/j.1365-2486.2011.02588.x
- Peterson, T. J., Saft, M., Peel, M. C., & John, A. (2021). Watersheds may not recover from drought. *Science*, 372(6543), 745-749. DOI: 10.1126/science.abd5085
- Pham-Duc, B., Sylvestre, F., Papa, F., Frappart, F., Bouchez, C., & Cretaux, J. F. (2020). The Lake Chad hydrology under current climate change. *Scientific Reports*, 10(1), 5498. DOI: 10.1038/s41598-020-62417-w
- Pimm, S. L. (1984). The complexity and stability of ecosystems. *Nature*, 307(5949), 321-326. DOI: 10.1038/307321a0
- Porkka, M., Virkki, V., Wang-Erlandsson, L., Gerten, D., Gleeson, T., Mohan, C.,Kummu, M. (2024). Notable shifts beyond pre-industrial streamflow and soil moisture conditions transgress the planetary boundary for freshwater change. *Nature Water*, 2(3), 262-273. DOI: 10.1038/s44221-024-00208-7
- Portela, A. P., Durance, I., Vieira, C., & Honrado, J. (2023a). Environmental filtering and environmental stress shape regional patterns of

riparian community assembly and functional diversity. *Freshwater Biology*, 68(8), 1428-1441. DOI: 10.1111/fwb.14138

- Portela, A. P., Durance, I., Vieira, C., & Honrado, J. (2023b). Response-effect trait overlap and correlation in riparian plant communities suggests sensitivity of ecosystem functioning and services to environmental change. *Science* of the Total Environment, 860, 160549. DOI: 10.1016/j.scitotenv.2022.160549
- Portela, A. P., Gonçalves, J. F., Durance, I., Vieira, C., & Honrado, J. (2023). Riparian forest response to extreme drought is influenced by climatic context and canopy structure. *Science* of the Total Environment, 881, 163128. DOI: 10.1016/j.scitotenv.2023.163128
- Pott, A., & Pott, V. J. (2004). Features and conservation of the Brazilian Pantanal wetland. *Wetlands Ecology and Management*, 12(6), 547-552. DOI: 10.1007/s11273-005-1754-1
- Rebelo, L.-M., McCartney, M. P., & Finlayson, M. C. (2011). The application of geospatial analyses to support an integrated study into the ecological character and sustainable use of Lake Chilwa. *Journal of Great Lakes Research*, 37, 83-92. DOI: 10.1016/j.jglr.2010.05.004
- Renaut, R. W., & Owen, R. B. (2023). Lake Naivasha. In The Kenya Rift Lakes: Modern and Ancient: *Limnology and Limnogeology* of Tropical Lakes in a Continental Rift (pp. 417-461). Springer Berlin Heidelberg. DOI: 10.1007/978-3-642-25055-2_11
- Richardson, K., Steffen, W., Lucht, W., Bendtsen, J., Cornell, S. E., Donges, J. F., ...Rockström, J. (2023). Earth beyond six of nine planetary boundaries. *Science Advances*, 9(37), eadh2458. DOI: 10.1126/sciadv.adh2458
- Ricketts, T. H., Watson, K. B., Koh, I., Ellis, A. M., Nicholson, C. C., Posner, S., Richardson, L. L., & Sonter, L. J. (2016). Disaggregating the evidence linking biodiversity and ecosystem services [Article]. *Nature Communications*, 7, 13106. DOI: 10.1038/ncomms13106
- Riebe, K., & Dressel, A. (2021). The impact on food security of a shrinking Lake Chad. *Journal of Arid Environments*, 189. DOI: 10.1016/j.jaridenv.2021.104486
- Riis, T., Kelly-Quinn, M., Aguiar, F. C., Manolaki, P., Bruno, D., Dufour, S. (2020). Global

Overview of Ecosystem Services Provided by Riparian Vegetation. *Bioscience*, 70(6), 501-514. DOI: 10.1093/biosci/biaa041

- Rogers, J. B., Stein, E. D., Beck, M. W., & Ambrose, R. F. (2020). The impact of climate change induced alterations of streamflow and stream temperature on the distribution of riparian species. *PloS one*, 15(11), e0242682. DOI: 10.1371/journal.pone.0242682
- Runge, K., Tucker, M., Crowther, T. W., Fournier de Lauriere, C., Guirado, E., Bialic-Murphy, L., & Berdugo, M. (2025). Monitoring Terrestrial Ecosystem Resilience Using Earth Observation Data: Identifying Consensus and Limitations Across Metrics. *Global Change Biology*, 31(3), e70115. DOI: 10.1111/ gcb.70115
- Seena, S., Baschien, C., Barros, J., Sridhar, K. R., Graça, M. A. S., Mykrä, H., & Bundschuh, M. (2022). Ecosystem services provided by fungi in freshwaters: a wake-up call. *Hydrobiologia*, 850(12-13), 2779-2794. DOI: 10.1007/ s10750-022-05030-4
- Small, N., Munday, M., & Durance, I. (2017). The challenge of valuing ecosystem services that have no material benefits. *Global Envi*ronmental Change-Human and Policy Dimensions, 44, 57-67. DOI: 10.1016/j.gloenvcha.2017.03.005
- Standish, R. J., Hobbs, R. J., Mayfield, M. M., Bestelmeyer, B. T., Suding, K. N., Battaglia, L. L., Thomas, P. A. (2014). Resilience in ecology: Abstraction, distraction, or where the action is? *Biological Conservation*, 177, 43-51. DOI: 10.1016/j.biocon.2014.06.008
- Sterner, R. W., Keeler, B., Polasky, S., Poudel, R., Rhude, K., & Rogers, M. (2020). Ecosystem services of Earth's largest freshwater lakes. *Ecosystem Services*, 41. DOI: 10.1016/j.ecoser.2019.101046
- Strickland, M. K., Jenkins, M. A., Ma, Z., & Murray, B. D. (2024). How has the concept of resilience been applied in research across forest regions? *Frontiers in Ecology and the Environment*, 22(3), e2703. DOI: 10.1002/ fee.2703
- Stromberg, J. C., McCluney, K. E., Dixon, M. D., & Meixner, T. (2012). Dryland Riparian Ecosystems in the American Southwest: Sen-

sitivity and Resilience to Climatic Extremes. *Ecosystems*, 16(3), 411-415. DOI: 10.1007/s10021-012-9606-3

- Thielen, D., Schuchmann, K.-L., Ramoni-Perazzi, P., Marquez, M., Rojas, W., Quintero, J. I., & Marques, M. I. (2020). Quo vadis Pantanal? Expected precipitation extremes and drought dynamics from changing sea surface temperature. *PloS one*, 15(1), e0227437. DOI: 10.1371/journal.pone.0227437
- Thomaz, S. M. (2023). Ecosystem services provided by freshwater macrophytes [Review]. *Hydrobiologia*, 850(12-13), 2757-2777. DOI: 10.1007/s10750-021-04739-y
- Tomas, W. M., Andrade, M. H., Berlinck, C. N., Bolzan, F., Camilo, A. R., Catella, A. C.,... Urbanetz, C. (2024). Eight basic principles for the elaboration of public policies and development projects for the Pantanal. *Conservation Science and Practice*. DOI: 10.1111/ csp2.13207
- Tomas, W. M., Berlinck, C. N., Chiaravalloti, R. M., Faggioni, G. P., Strussmann, C., Libonati, R.,...Morato, R. (2021). Distance sampling surveys reveal 17 million vertebrates directly killed by the 2020's wildfires in the Pantanal, Brazil. *Scientific Reports*, 11(1), 23547. DOI: 10.1038/s41598-021-02844-5
- Tomas, W. M., de Oliveira Roque, F., Morato, R. G., Medici, P. E., Chiaravalloti, R. M., Tortato, F. R., Penha, J. M. F.,...Junk, W. J. (2019).
 Sustainability Agenda for the Pantanal Wetland: Perspectives on a Collaborative Interface for Science, Policy, and Decision-Making. Tropical Conservation Science, 12. DOI: 10.1177/1940082919872634
- United Nations. (2022). Sustainable Development Goals. Retrieved 2022/06/01 from https:// sdgs.un.org/goals
- UNESCO. (2025). Pantanal Conservation Area. https://whc.unesco.org/en/list/999/
- Van der Ploeg, S., De Groot, R. S., & Wang, Y. (2010). The TEEB Valuation Database: overview of structure, data and results. Foundation for sustainable development, Wageningen, the Netherlands, 1-247.
- van Dijk, A. I. J. M., Beck, H. E., Crosbie, R. S., de Jeu, R. A. M., Liu, Y. Y., Podger, G. M., Timbal, B., & Viney, N. R. (2013). The

Millennium Drought in southeast Australia (2001–2009): Natural and human causes and implications for water resources, ecosystems, economy, and society. *Water Resources Research*, 49(2), 1040-1057. DOI: 10.1002/wrcr.20123

- Van Meerbeek, K., Jucker, T., & Svenning, J. C. (2021). Unifying the concepts of stability and resilience in ecology. *Journal of Ecolo*gy, 109(9), 3114-3132. DOI: 10.1111/1365-2745.13651
- Wantzen, K. M., Assine, M. L., Bortolotto, I. M., Calheiros, D. F., Campos, Z., Catella, A. C., ...Urbanetz, C. (2024). The end of an entire biome? World's largest wetland, the Pantanal, is menaced by the Hidrovia project which is uncertain to sustainably sup-

port large-scale navigation. *Science of the Total Environment*, 908. DOI: 10.1016/j. scitotenv.2023.167751

- Xu, X., Chen, M., Yang, G., Jiang, B., & Zhang, J. (2020). Wetland ecosystem services research: A critical review. *Global Ecology* and Conservation, 22. DOI: 10.1016/j.gecco.2020.e01027
- Zieritz, A., Sousa, R., Aldridge, D. C., Douda, K., Esteves, E., Ferreira-Rodriguez, N., Mageroy, J. H., Nizzoli, D., Osterling, M., Reis, J., Riccardi, N., Daill, D., Gumpinger, C., & Vaz, A. S. (2022). A global synthesis of ecosystem services provided and disrupted by freshwater bivalve molluscs. *Biological Reviews Cambridge Philosophical Society*, 97(5), 1967-1998. DOI: 10.1111/brv.12878